

Papers on combinatorial chemistry or solid-phase synthesis from other journals – May 2001

A novel concept in combinatorial chemistry in solution with the advantages of solid-phase synthesis: formation of N-betaines by multicomponent domino reactions. Tietze, L.F.; Evers, H.; Topken, E. *Angew. Chem.*, (2001), 40(5), 903-905.

Revolutionizing resin handling for combinatorial synthesis. Atrash, B.; Bradley, M.; Kobylecki, R.; Cowell, D.; Reader, J. *Angew. Chem.*, (2001), 40(5), 938-941.

Tracelessness unmasked: a general linker nomenclature. Comely, A.C.; Gibson, S.E. *Angew. Chem.*, (2001), 40(6), 1012-1032.

A structurally biased combinatorial approach for discovering new anti-picornaviral compounds. Tsang, S.K.; Cheh, J.; Isaacs, L.; Joseph-McCarthy, D.; Choi, S-K.; Pevear, D.C.; Whitesides, G.M.; Hogle, J.M. *Chem. Biol.* (2001), 8(1), 33-45.

An enzyme-labile safety catch linker for synthesis on a soluble polymeric support. Grether, U.; Waldmann, H. *Chem. Eur. J.* (2001), 7(5), 959-971.

Solid phase synthesis of pyridazine derivatives using polymer-bound sodium benzenesulfinate. Chen, Y.; Lam, Y.; Lee, S-Y. *Chem. Lett.* (2001), (3), 274-275.

Polymeric scavenger reagents in organic synthesis. Eames, J.; Watkinson, M. *Eur. J. Org. Chem.* (2001), (7), 1213-1224.

Quantitative determination of resin loading in solid-phase organic synthesis using ¹³C MAS NMR. Hany, R.; Rentsch, D.; Dhanapal, B.; Obrecht, D. *J. Comb. Chem.* (2001), 3(1), 85-89.

Convenient preparation of 4-formyl-3,5-dimethoxyphenol and its incorporation into linkers and resins for solid-phase synthesis. Jin, J.; Graybill, T.L.; Wang, M.A.; Davis, L.D.; Moore, M.L. *J. Comb. Chem.* (2001), 3(1), 97-101.

A rigid linker-scaffold for solid-phase synthesis of dimeric pharmacophores. Pattarawarapan, M.; Chen, J.; Steffensen, M.; Burgess, K. *J. Comb. Chem.* (2001), 3(1), 102-116.

Williamson ether synthesis on solid support: substitution versus elimination. Weissberg, A.; Dahan, A.; Portnoy, M. *J. Comb. Chem.* (2001), 3(2), 154-156.

The Preparation of a New "Safety Catch" Ester Linker for Solid-Phase Synthesis. Beech, C.L.; Coope, J.F.; Fairley, G.; Gilbert, P.S.; Main, B.G.; Ple, K. *J. Org. Chem.* (2001), 66(7), 2240-2245.

(Cyanomethyl)trialkylphosphonium iodides: efficient reagents for the intermolecular

alkylation of amines with alcohols in solution and on solid phase. Zaragoza, F.; Stephensen, H. *J. Org. Chem.* (2001), 66(7), 2518-2521.

A modular approach for the synthesis of oligosaccharide mimetics. Patel, A.; Lindhorst, T.K. *J. Org. Chem.* (2001), 66(8), 2674-2680.

Solid-phase synthesis of β -lactams via the Miller hydroxamate approach. Meloni, M.M.; Taddei, M. *Org. Lett.* (2001), 3(3), 337-340.

Potent integrin antagonists from a small library of RGD-including cyclic pseudopeptides. Belvisi, L.; Bernardi, A.; Checchia, A.; Manzoni, L.; Potenza, D.; Scolastico, C.; Castorina, M.; Cupelli, A.; Giannini, G.; Carminati, P.; Pisano, C. *Org. Lett.* (2001), 3(7), 1001-1004.

The H-phosphonate approach to the synthesis of phosphopeptides on solid phase. Kupihar, Z.; Kele, Z.; Toth, G.K. *Org. Lett.* (2001), 3(7), 1033-1035.

Fmoc solid phase synthesis of polyamides containing pyrrole and imidazole amino acids. Wurtz, N.R.; Turner, J.M.; Baird, E.E.; Dervan, P.B. *Org. Lett.* (2001), 3(8), 1201-1203.

High-resolution analysis of a 144-membered pyrazole library from combinatorial solid phase synthesis by using electrospray ionisation Fourier transform ion cyclotron resonance mass spectrometry. Schmid, D.G.; Grosche, P.; Jung, G. *Rapid Commun. Mass Spectrom.* (2001), 15(5), 341-347.

Solid-phase synthesis of quinazoline-2,4-diones using S_NAr reaction. Makino, S.; Suzuki, N.; Nakanishi, E.; Tsuji, T. *Synlett* (2001), (3), 333-336.

Solid-phase free radical addition of thiols to allyl ethers. Plourde, R., Jr.; Johnson, L.L., Jr.; Longo, R.K. *Synlett* (2001), (3), 439-441.